

HOSPITAL EPISODES DUE TO ANTIDEPRESSANT OVERDOSE: AN UNDER-UTILISED SOURCE OF PHARMACOVIGILANCE DATA

Cooke MJ¹ ; Tomlinson D² ; Kitching G³ ; Waring WS⁴

1. Pharmacy Department, York Hospital, York, UK
2. Medicines Management, North Yorkshire & York Primary Care Trust, York, UK
3. Emergency Department, York Hospital, York, UK
4. Acute Medical Unit, York Hospital, York, UK

ABSTRACT

Background: Antidepressant agents are commonly implicated in drug overdose, and the toxicological profile varies between agents. Clinical data concerning overdoses are not systematically sought or evaluated in pharmacovigilance. The present study sought to examine the feasibility of collecting Emergency Department data concerning antidepressant overdose.

Methods : Presentations to York Hospital due to intentional antidepressant overdose were studied between 2010 and 2011. Data collected were the type of antidepressant, dose, co-ingested drugs, duration of hospital stay, and need for critical care. Community National Health Service prescription data were evaluated across York and North Yorkshire region.

Results : There were 250 overdose episodes. These involved a selective serotonin reuptake inhibitor (SSRI) in 183 (73.2%), and a tricyclic in 45 (18.0%), equivalent to 24 episodes per 100,000 prescription items (95% CI 21-28 per 100,000) and 11 per 100,000 (8-15 per 100,000) respectively (P<0.0001). Citalopram was the most commonly prescribed, and associated with 22 overdose episodes per 100,000 (17-27 per 100,000). Fluoxetine was associated with 32 overdose episodes per 100,000 (24-41 per 100,000) (P=0.032 versus citalopram), whereas the lower rates were observed for amitriptyline (13, 9-17 per 100,000) (P=0.004) and dosulepin (2, 0-10 per 100,000) (P=0.001).

Conclusions : A higher than expected number of overdose episodes involved an SSRI based on National Health Service primary care prescribing, and fewer episodes involved a tricyclic antidepressant. Clinical outcomes differed between agents, indicating the feasibility of using Emergency Department data to detect different patterns of toxicity between antidepressants. Further work is required to examine whether systematic collection of clinical toxicology data might enhance existing pharmacovigilance systems.

Corresponding author: WS Waring ; Acute Medical Unit , York Teaching Hospital NHS Foundation Trust , Wigginton Road , York , YO31 8HE ; **Telephone:** 01904 726276 ; **E-mail:** stephen.waring@york.nhs.uk

Citation: Cooke MJ, Tomlinson D, Kitching G, Waring WS (2013) Hospital Episodes Due to Antidepressant Overdose: An Under-Utilised Source of Pharmacovigilance Data. Journal of Advanced Pharmaceutical Science And Technology - 1(2):16-21. <https://doi.org/10.14302/issn.2328-0182.japst-13-185>

Running title: Antidepressant prescribing versus overdose

Keywords: Antidepressant; drug safety; overdose; pharmacovigilance; prescribing patterns; SSRI; toxicology; tricyclic antidepressant

Academic Editor: Satheesh Podaralla, Research Scientist : Formulation R&D SRI International Menlo Park, CA

INTRODUCTION

Intentional drug overdose is an important cause of Emergency Department attendances in the United Kingdom and elsewhere [1]. Antidepressant agents are commonly implicated in drug overdose, and may result in significant morbidity and mortality [2,3]. Clinical outcomes after intentional antidepressant overdose vary substantially depending on the type of agent involved and dose ingested. For example, monoamine oxidase inhibitor (MAOI) ingestion can result in accumulation of norepinephrine within the central nervous system, and may cause profound haemodynamic instability, seizures, coma and death [4,5]. Tricyclic antidepressants are capable of blocking sodium channel conductance such that overdose is characterised by the occurrence of life-threatening arrhythmia, severe hypotension, generalised seizures, coma, and death [6,7]. In contrast, overdose involving a selective serotonin reuptake inhibitor (SSRI) is characterised by less overall toxicity, fewer life-threatening complications, and a lower mortality than that attributed to MAOI or tricyclic antidepressants [8-10].

In recent years, there has been a trend towards increase antidepressant prescribing due widened clinical indications for pharmacological treatment. Prescribing patterns have changed such that MAOIs and tricyclic antidepressants are less widely relied upon, and there has been a substantial increase in the use of SSRIs and novel agents including duloxetine, mirtazapine, and venlafaxine [11,12]. These changes have been prompted by several factors, recommendations in clinical guidelines, adverse effect profile of specific drugs, marketing activity by the pharmaceutical industry, patient preference, and the perceived risk of self-harm in individual patients. Many existing pharmacovigilance systems have been designed to allow collection of data concerning medications used in accordance with the Marketing Authorisation, namely adverse effects arising from intended therapeutic use including prescribed dosages. In contrast, data concerning adverse outcomes in the setting of intentional drug overdose is not collected in a systematic manner, and these data tend to emerge as individual case reports, published case series, and abstracts at scientific meetings. Comparatively few data are collected systematically in this patient group, and there is a need for a better understanding of the link between exposure and extent of exposure to particular drugs and clinical outcomes. The present study sought to examine the possibility of using Emergency Department data to identify patterns of antidepressant overdose with respect to prevailing prescribing practices, and to evaluate clinical outcomes between specific agents as a measure of toxicity.

METHODS

Clinical data

York Hospital serves a semi-rural population of around 350,000 people in York and the North Yorkshire region, and receives approximately 70,000 Emergency Department attendances annually. This includes almost 900 attendances due to intentional drug overdose (1.3%) [13]. Adults that present after drug overdose may be discharged home after completion of medical and psychiatric assessment, or admitted to the Acute Medical Unit if ongoing medical treatment is required or

the patient is too drowsy or intoxicated to allow detailed psychiatric assessment. Patients may be admitted to a critical care area if there is severe haemodynamic disturbance or if non-invasive or invasive ventilatory support may be required. The study population consisted of all patients aged ≥ 16 years that presented to the Emergency Department due to drug overdose between January 2010 and December 2011 inclusive. Data collected were age, gender, date and time of ingestion, type of agent ingested, dose, co-ingested drugs and ethanol, admission to hospital, duration of hospital stay, and if transfer to a critical care area was required.

Prescribing data

National Health Service primary care prescribing data in York and North Yorkshire region are routinely collected and examined for the purposes of evaluating prescribing trends and variance by prescribers across a wide range of therapeutic areas. The database was interrogated for all prescriptions items for antidepressant agents in 2010 and 2011. Antidepressants were classified as a tricyclic, SSRI, or other antidepressant in accordance with the categories listed in the British National Formulary [14].

Data Analyses

Data concerning specific antidepressant agents was considered as a proportion of all overdose episodes or prescribing items. Overdose episodes were expressed per 100,000 prescription items, and comparison made between agents using Chi square proportional tests. The ingested dose was expressed as a multiple of the World Health Organization 'Defined Daily Dose' (DDD) to allow comparison between agents [15]. Analyses were performed using MedCalc statistical software v.9.1.0.1 (MedCalc, Mariakerke, Belgium), and P-values < 0.05 were accepted as statistically significant [16].

RESULTS

There were 1092 Emergency Department attendances due to intentional drug overdose, including 250 episodes that involved an antidepressant (22.9%). The median age was 37 years (population range 16-85 years), and there were 165 women (66%) (Table 1). Hospital presentations involved an SSRI in 183 (73.2%), a TCA in 45 (18.0%), and another antidepressant in 22 (8.8%). Patients that ingested a TCA were more likely to require admission to hospital, had a longer duration of hospital stay, and were more likely to require transfer to a critical care area (Table 1). One death occurred in a 34-year-old woman with a past history of chronic alcohol excess that presented to hospital due to asystolic cardiac arrest after overdose involving unknown quantities of escitalopram, paracetamol, and codeine. Patterns of antidepressant overdose in men and women are presented in Table 2.

There were 1,304,795 prescription items for antidepressants in the York and North Yorkshire region,

(Continued on page 18)

and these included 749,144 (57.4%) items for a SSRI, 427,127 (32.7%) for a TCA, and 129,243 (9.9%) others. The overall occurrence of antidepressant overdose was equivalent to 19 episodes per 100,000 prescription items (95% CI 17-22 per 100,000 scripts). A higher number of overdoses were observed in the SSRI group compared to the TCA group: 24 (95% CI 21-28) versus 11 (95% CI 8-15) respectively ($P < 0.0001$). Citalopram was the most widely prescribed, and used as a reference comparator (Table 3). Fluoxetine was associated with a higher number of overdose episodes: 32 per 100,000 scripts (24-41) versus 22 (17-27) respectively ($P = 0.032$). Fewer than expected overdose episodes involved amitriptyline (13, 95% CI 9-17, $P = 0.004$) and dosulepin (2, 95% CI 0-10, $P = 0.001$) (Figure 1).

DISCUSSION

TABLE 1. Summary characteristics of patients presenting after intentional antidepressant overdose presented as median (interquartile range) and proportions. * $P = 0.0082$, ** $P = 0.0024$ by Mann Whitney tests, and † $P = 0.0181$, and †† $P = 0.0002$ by Chi square proportional tests versus SSRI group.

	SSRI n=183	Tricyclic Anti- depressant n=45	Other Anti- depressants n=22
Women	132 (72.1%)	24 (53.3%)	9 (40.9%)
Men	51 (27.9%)	21 (46.7%)	13 (59.1%)
Age (years)	35 (22-47)	37 (30-44)	38 (30-47)
Alcohol co-ingestion	71 (38.8%)	22 (48.9%)	11 (50.0%)
Other drug co-ingestion	127 (69.4%)	33 (73.3%)	19 (86.4%)
Drug dose as DDD multiple	14 (3-31)	7 (0-22)**	15 (0-37)
Hospital admission	102 (55.7%)	39 (86.7%)†	15 (68.2%)
Critical care admission	3 (1.6%)	7 (15.6%)††	0 (0.0%)
Duration of hospital stay (days)	1 (1-1)	2 (1-4)*	1 (1-2)


TABLE 2. Comparison of antidepressant overdose between men and women. * $P = 0.0488$, ** $P = 0.0141$, *** $P = 0.0130$ by Chi square proportional tests.

	Women n=183	Men n=45
SSRI	132 (80.0%, 66.9-94.9%)	51 (60.0%, 44.7-78.9%)
Citalopram	57 (34.6%)	26 (30.6%)
Fluoxetine	38 (23.0%)	15 (17.7%)
Venlafaxine	15 (9.1%)	4 (4.7%)
Sertraline	15 (9.1%)	4 (4.7%)
Escitalopram	4 (2.4%)	0 (0.0%)
Paroxetine	3 (1.8%)	0 (0.0%)
Tricyclics	24 (14.6%, 9.3-21.6%)	21 (24.7%, 15.3-37.8%)
Amitriptyline	22 (13.3%)	17 (20.0%)
Lofepramine	1 (0.6%)	1 (1.2%)
Clomipramine	1 (0.6%)	0 (0.0%)
Dosulepin	0 (0.0%)	2 (2.4%)*
Imipramine	0 (0.0%)	1 (1.2%)
Others	9 (5.5%, 2.5-10.4%)	13 (15.3%, 8.1-26.2%)*
Mirtazapine	8 (4.8%)	12 (14.1%)**
Duloxetine	1 (0.6%)	0 (0.0%)
Trazodone	0 (0.0%)	1 (1.2%)

Around five and a half thousand deaths were attributed to drug poisoning in England and Wales alone in 2010-2011, and antidepressants account for a high proportion of these. The age-standardised annual mortality rates attributed to antidepressant overdose are 7.1 (95% CI 6.1-8.1) and 6.7 (95% CI 5.8-7.6) per million in men and women respectively, which are almost double those attributed to paracetamol poisoning of 3.1 (2.4-3.7) and 3.6 (3.0-4.3) deaths per million in men and women respectively [17]. Within these deaths, differences exist between agents as a result of variations in their accessibility and inherent toxicity [18]. Citalopram was the most widely prescribed antidepressant and most likely to be implicated in overdose, despite mounting data that it is capable of prolonging the QT interval and may predispose to arrhythmia [19, 20]. The fatal toxicity index has been proposed as a measure that takes account of the number of prescription items and, for example [21]. This indicates between 11 and 35 deaths per million antidepressant prescription items, including 27 to 47 deaths per million amitriptyline prescription items [22, 23]. An alternative approach has been to consider the case fatality, expressed as a ratio of fatal to non-fatal self-poisonings; the case fatality has been reported for tricyclic antidepressants and SSRIs as 13.8 (95% CI 13.0-14.7) and 0.5 (95% CI 0.4-0.7) respectively ($P < 0.0001$) [24]. Only one death was

(Continued on page 19)

FIGURE 1. The proportion of overdose episodes related to specific antidepressant agents, expressed as percentage of all episodes with 95% confidence intervals; the corresponding community prescriptions are expressed as a percentage of all antidepressant prescription items.



observed in the present study, equivalent to 0.8 deaths per million prescription items (95% CI 0.0-4.3 deaths per million), indicating that only a very small proportion of antidepressant poisoning deaths occur in hospital.

As expected, the present findings indicate that overdose involving tricyclics was associated with a greater need for hospital admission, longer hospital stay, and greater need for critical care compared to overdose involving SSRI or other antidepressants. These observations may be explained as a result of differing intrinsic toxicity because the ingested dose was comparatively smaller in the tricyclic group, and there were no significant differences in co-ingested drugs and ethanol [25]. Previous reports indicate that intentional overdose involving MAOI or tricyclic antidepressants may result in severe arrhythmia and prolonged generalised seizures that are resistant to conventional therapy [26]. In contrast, overdose involving SSRIs may provoke dose-dependent generalised seizures in around 10%, and these are usually self-limiting or promptly terminated by benzodiazepine administration [27]. SSRI poisoning gives rise to prolongation of the QT interval in around 5% of patients, but the occurrence of life-threatening cardiac arrhythmia is thought to be very rare [28, 29].

TABLE 3. Antidepressant overdose episodes and community prescription items in 2010-2011. *P=0.033, **P=0.005 by Chi square proportional tests versus citalopram group; †P<0.0001 versus SSRI group.

Antidepressant	Prescription items	Overdose episodes	Episodes per 100,000 scripts	Likelihood ratio	95% CI
SSRIs					
Citalopram	380640	83	22 (17-27)	1.00	-
Escitalopram	27393	4	15 (4-37)	0.67	0.25-1.80
Fluoxetine	167096	53	32 (24-41)	1.45*	1.03-1.22
Paroxetine	35479	3	9 (2-25)	0.39	0.12-1.22
Sertraline	79807	21	26 (16-40)	1.21	0.75-1.95
Venlafaxine	57784	19	33 (20-51)	1.51	0.92-2.48
All SSRIs	748199	183	24 (21-28)		
TCA's					
Amitriptyline	310123	39	13 (9-17)	0.58**	0.39-0.84
Clomipramine	11047	1	9 (0-5)	0.42	0.06-2.98
Dosulepin	69579	2	3 (0-10)	0.13**	0.03-0.54
Imipramine	6321	1	16 (0-9)	0.73	0.10-5.21
Lofepramine	15042	2	13 (2-5)	0.61	0.15-2.48
All TCA's	412112	45	11 (8-15)†		
Others					
Duloxetine	9883	1	10 (0-56)	0.46	0.06-3.33
Mirtazapine	88705	20	23 (14-35)	1.03	0.63-1.68
Trazodone	28729	1	4 (1-19)	0.16	0.02-1.15
All others	127317	22	17 (11-26)		
All antidepressants	1287628	250	19 (17-22)		

The study design did not directly address the factors that might give rise to preferential involvement of particular drugs amongst patients that attend hospital after intentional overdose, and a number of possible explanations exist. Firstly, a high level of awareness of the comparative toxicity of various antidepressant agents and judicious prescribing would result in greater reliance upon SSRI agents, and less reliance upon tricyclic or MAOI agents when issuing prescriptions to patients that are considered to be at increased risk of self-harm. Secondly, the tendency towards increasing antidepressant use coupled with SSRIs being relied upon as

first-line treatment means that a greater number of 'low risk' patients will receive an SSRI, and this group may be more likely to seek medical attention after intentional overdose. Thirdly, tricyclic antidepressants may be prescribed where SSRI treatment has failed, demarcating a subset of patients with more resistant or severe depressive illness that may be less inclined to seek medical attention after overdose. Fourthly, tricyclic antidepressants may have been initiated in the past, and used to maintain long-term remission of symptoms, and this patient subgroup may be less likely to exhibit self-harm behaviour. Lastly, tricyclics may be more likely to be used for disorders such as neuropathic pain, where the risk of intentional overdose is expected to be lower.

CONCLUSIONS

SSRIs were more highly represented amongst overdose patients than might have been predicted from the numbers of prescription items, whereas fewer episodes involved a tricyclic antidepressant. These data suggest a preferential use of SSRI rather than tricyclic antidepressants in patients at risk of self-harm. Further work is required to understand the factors that influence antidepressant prescribing so as to minimise the risk of harm in this at-risk patient population. These findings indicate that Emergency Department data are sufficiently sensitive to detect different clinical outcomes between antidepressant agents in overdose. These data may make a useful addition to the existing sources that are relied upon in conventional pharmacovigilance programmes. Further work is required to explore the feasibility of systematic clinical data collection and how these data might be used to inform prescribers in order to minimise adverse outcomes in patients at high risk of self-harm.

ACKNOWLEDGEMENTS

We would like to thank Jackie Snowden, York Hospital for assistance with retrieval of the case notes to enable data collection and analyses.

REFERENCES

1. Waring, W.S., and McGettigan, P. (2011) Clinical toxicology and drug regulation: a United Kingdom perspective. *Clin. Toxicol.* 49, 452-456
2. Thanacoody, H.K., and Thomas, S.H. (2003) Antidepressant poisoning. *Clin. Med.* 3, 114-118
3. Waring, W.S. (2012) Clinical use of antidepressant therapy and associated cardiovascular risk. *Drug Healthc. Patient Saf.* 4, 93-101
4. Sarko, J. (2000) Antidepressants, old and new. A review of their adverse effects and toxicity in overdose. *Emerg. Med. Clin. North Am.* 18, 637-654
5. Thanacoody, H.K., and Waring, W.S. (2008) Toxins that affect the cardiovascular system. *Clin. Med.* 8, 92-95
6. Koegelenberg, C.F., Joubert, Z.J., and Irusen, E.M. (2012) Tricyclic antidepressant overdose necessitating ICU admission. *S. Afr. Med. J.* 102, 293-294
7. Levine, M., Brooks, D.E., Franken, A., and Graham, R. (2012) Delayed-onset seizure and cardiac arrest after amitriptyline overdose, treated with intravenous lipid emulsion therapy. *Pediatrics.* 130, e432-e438
8. Isbister, G.K., Bowe, S.J., Dawson, A., and Whyte, I.M. (2004) Relative toxicity of selective serotonin reuptake inhibitors (SSRIs) in overdose. *J. Toxicol. Clin. Toxicol.* 42, 277-285
9. Kelly, C.A., Dhaun, N., Laing, W.J., Strachan, F.E., Good, A.M. et al. (2004) Comparative toxicity of citalopram and the newer antidepressants after overdose. *J. Toxicol. Clin. Toxicol.* 42, 67-71
10. Flanagan, R.J. (2008) Fatal toxicity of drugs used in psychiatry. *Hum. Psychopharmacol.* 23(Suppl 1), 43-51
11. Wong, A., Taylor, D.M., Ashby, K., Robinson, J. (2010) Changing epidemiology of intentional antidepressant drug overdose in Victoria, Australia. *Aust. N. Z. J. Psychiatry.* 44, 759-764
12. Bergen, H., Murphy, E., Cooper, J., Kapur, N., Stalker, C. et al. (2010) A comparative study of non-fatal self-poisoning with antidepressants relative to prescribing in three centres in England. *J. Affect. Disord.* 123, 95-101
13. James, N.J., Hussain, R., Moonie, A., Richardson, D., and Waring, W.S. (2012) Patterns of admissions in an acute medical unit: priorities for service development and education. *Acute Med.* 11, 74-80
14. Joint Formulary Committee. British National Formulary (online) London: BMJ Group and Pharmaceutical Press <http://www.medicinescomplete.com> Accessed on 05th December 2012
15. Armstrong, T.M., Davies, M.S., Kitching, G., and Waring, W.S. (2012) Comparative drug dose and drug combinations in patients that present to hospital due to self-poisoning. *Basic Clin. Pharmacol. Toxicol.* 111, 356-360
16. Schoonjans, F., Zalata, A., Depuydt, C.E., and Comhaire, F.H. (1995) MedCalc: a new computer program for medical statistics. *Comput. Methods Programs Biomed.* 48, 257-262
17. The Office for National Statistics (ONS). Deaths related to drug poisoning in England and Wales, 2011. Statistical Bulletin 29/08/2012. <http://www.ons.gov.uk/ons/index.html> Accessed on 05th December 2012
18. White, N., Litovitz, T., and Clancy, C. (2008) Suicidal antidepressant overdoses: a comparative analysis by antidepressant type. *J. Med. Toxicol.* 4, 238-250
19. Friberg, L.E., Isbister, G.K., and Duffull, S.B. (2006) Pharmacokinetic-pharmacodynamic modelling of QT interval prolongation following citalopram overdoses. *Br. J. Clin. Pharmacol.* 61, 177-190
20. Cooke, M.J., and Waring, W.S. (2012) Citalopram and cardiac toxicity. *Eur. J. Clin. Pharmacol.* 69, 755-760.
21. Rose, J.C., and Unis, A.S. (2000) A mortality index for postmarketing surveillance of new medications. *Am. J. Emerg. Med.* 18, 176-179
22. Henry, J.A., and Antao, C.A. (1992) Suicide and fatal antidepressant poisoning. *Eur. J. Med.* 1, 343-348

23. Buckley, N.A., and McManus, P.R. (1998) Can the fatal toxicity of antidepressant drugs be predicted with pharmacological and toxicological data. *Drug Saf. 18*, 369-381
24. Hawton, K., Bergen, H., Simkin, S., Cooper, J., Waters, K. et al. (2010) Toxicity of antidepressants: rates of suicide relative to prescribing and non-fatal overdose. *Br. J. Psychiatry. 196*, 354-358
25. Koski, A., Vuori, E., and Ojanperä, I. (2005) Newer antidepressants: evaluation of fatal toxicity index and interaction with alcohol based on Finnish postmortem data. *Int. J. Legal Med. 119*, 344-348
26. Waring, W.S., and Wallace, W.A. (2007) Acute myocarditis after massive phenelzine overdose. *Eur. J. Clin. Pharmacol. 63*, 1007-1009
27. Waring, W.S., Gray, J.A., and Graham, A. (2008) Predictive factors for generalized seizures after deliberate citalopram overdose. *Br. J. Clin. Pharmacol. 66*, 861-865
28. Tarabar, A.F., Hoffman, R.S., and Nelson, L. (2008) Citalopram overdose: late presentation of torsades de pointes (TdP) with cardiac arrest. *J. Med. Toxicol. 4*, 101-105
29. Waring, W.S., Graham, A., Gray, J., Wilson, A.D., Howell, C., et al. (2010) Evaluation of a QT nomogram for risk assessment after antidepressant overdose. *Br. J. Clin. Pharmacol. 70*, 881-885