

# Dose Physiologically Performing Vigorous Exercises Good for Body Over Time

Nasim Habibzadeh<sup>1,\*</sup>

<sup>1</sup>PhD in Sport Science, Department of Sport Science, Teesside University, UK

## Introduction

The body physiological responses to the exercise paradigms rely on the exercise volume (or pace rate) and exercise duration. During exercise the glycolysis, fat metabolisms and oxygen consumption increase to meet exercise demands that is accompanied with enhanced cardiac and respiratory systems workouts. The body temperature is also dramatically will raise. Depending on the exercise intensity rates, there are some physical stress on bones and musculature systems [1].

Performing exercises (i.e walking, running, cycling) in higher level of volume may increase the body stamina and improve the overall fitness. In this relation, several studies showed that various exercises can lead to faster workouts in daily tasks as the body segments and systems depending

on the individual unique capacities can adapt to the different physiological conditions. For example, it has been suggested that executing vigorous exercise is good for disabled or elderly populations as it makes them to be more active. It is believed that working on higher intensity of their abilities would cause this frail individual to perform more movements over time, although necessary cautions such as the amount of applied exercise intensity are required to take into account at these occasions [2,3].

Some other evidences recommended that vigorous exercise such as brisk walking can result in some health benefits such as enhancing weight loss, increasing more mechanical stress on muscle and bones to strengthen musculoskeletal system in spite of higher level of muscle soreness [4]. Although, generally not all individual can walk

**Corresponding author:** Nasim Habibzadeh, PhD in Sport Science, Department of Sport Science, Teesside University, UK, Email: [nasimhabibzadeh@yahoo.com](mailto:nasimhabibzadeh@yahoo.com)

**Citation:** Nasim Habibzadeh (2018) Dose Physiologically Performing Vigorous Exercises Good for Body Over Time. International Physiology Journal - 2(1):1-2. <https://doi.org/10.14302/issn.2578-8590.ipj-18-2544>

**Keywords:** Exercise

**Received:** Dec 13, 2018

**Accepted:** Dec 13, 2018

**Published:** Dec 14, 2018

too fast or are able to perform any intensive exercise, so that performing vigorous exercise cannot be developed to all people as different individuals have non-identical physical abilities in physiological terms. This is because the tolerated exercise volume (or speed rate) during various modes of exercises is different between individuals. Resultantly, every person has one unique peak exercise performance capability. Further, there are some kind of specific people such as patients with cardiovascular (CVD) difficulties who are not able to perform exercise in high level of intensities as their health status can be at more risks at that condition [5]. This type of people can manage their health condition with executing moderated kind of exercise modalities (i.e. mild walking at slow pace) to stay healthy with more care. Regular exercise like 10-15 min walking regimen per each day can be beneficial for CVD patients.

*More Importantly, the Question Which Can Come Up, is that what Really are the Side - Effect of Performing the Vigorous Exercises in Long-Term ?*

Some experts assume that the side-effects of vigorous exercise or over-training are pretty serious. Irritably, losing of appetite are primarily negative-effects of vigorous which can lead to the poor or inability to complete the related workouts or task performances. Other adverse clinical outcomes are fatigue, aching muscles and lethargy [6].

The constant vigorous exercise can affect bone frames. Intense exercise loads can increase strain on some particular bones such as back and in part spine areas and can cause risk of disk compression. Strenuous exercise also can increase an abnormal tilt of the pelvis or hip bones which can lead to the hip fractures and thus disabilities in final stage [7]. Although choosing the type of exercise in this regards is an important determinant.

However, the people competencies at different exercise domains are vary. Some people hereditary are born to work hard at different aspects and so there are always some exceptions in various areas such those high level elites as they have many different skills and expertise but not all people are the same [8,9]. Therefore, it seems performing moderate type of exercise is better choice for ordinary people since vigorous exercises are not suit for all. The simplest type

of moderate exercise is walking which can be performed in any locations without any specific physical or economical costs.

## References

1. Brooks GA, Fahey TD. Exercise Physiology—Human Biogenetics and its Applications.
2. Macmillan Publishing Company, 1985 Bruce B, Fries JF, Hubert H. Regular vigorous physical activity and disability development in healthy overweight and normal-weight seniors: a 13-year study. *Am J Public Health*. 2008;98:1294–1299
3. McPhee JS, French DP, Jackson D, Nazroo J, Pendleton N, Degens H. Physical activity in older age: perspectives for healthy ageing and frailty. *Biogerontology*. 2016 Jun;17(3):567-80
4. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC, Swain DP; American College of Sports Medicine. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011 Jul;43(7):1334-59
5. Sharman JE, La Gerche A, Coombes JS. Exercise and cardiovascular risk in patients with hypertension. *Am J Hypertens*. 2015 Feb;28(2):147-58
6. Deuk-Ja Oh, Hyeon-Ok Hong, Bo-Ae Lee. The effects of strenuous exercises on resting heart rate, blood pressure, and maximal oxygen uptake. *J Exerc Rehabil*. 2016 Feb; 12(1): 42– 46
7. Scherr J, Braun S, Schuster T, Hartmann C, Moehlenkamp S, Wolfarth B, Pressler A, Halle M. 72-h kinetics of high-sensitive troponin T and inflammatory markers after marathon. *Med Sci Sports Exerc*. 2011;43:1819 –1827
8. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med*. 2002;346:793–80
9. Iarizon E, Uy-Evanado A, Reinier K, Teodorescu C, Narayanan K, Jouven X, Gunson K, Jui J, Chugh SS. Sudden cardiac arrest during sports activity in middle age. *Circulation*. 2015;131:1384–1391