

The Pineal Hypothesis for Drug Dependence

Francesco Crespi^{1,*}

¹Biology, GSK Verona, Italy

Abstract

The pineal gland constitutes a major neuroendocrine organ in the brain. By mean of its neurohormone melatonin it transduces exogenous signals such as circadian and seasonal variations of light and temperature into proper hormonal changes which adjust and adapt internal endocrine functions. Alteration of circadian rhythms has been associated with affective disorders, psychosomatic diseases and cancer.

It has been observed that light deprivation, which stimulates (the enzymes responsible for) melatonin production in the pineal, enhances the animal's ethanol preference. Similarly, administration of the pineal hormone to rats maintained under normal conditions of constant photoperiod also induced ethanol drinking.

Our hypothesis is that in normal conditions melatonin might be acting as a cerebral "pacemaker", sensitive to endogenous as well as exogenous stimuli in the attempt to maintain an equilibrate circadian interaction between the cerebral activities of endogenous aminergic and opiates systems.

Abnormal states (i.e. drug abuse) could result in altered pineal activity, then in rhythmically altered functions of cerebral opiates and/or monoamine neurotransmitters. This may led to the development of a "reward - urge for drug rhythm" resulting in craving, ending in addiction.

Corresponding author: Francesco Crespi, Biology, GSK Verona, Italy. Email: fm.crespi@libero.it

Citation: Francesco Crespi (2018) The Pineal Hypothesis for Drug Dependence. International Physiology Journal - 2(1):3-8. <https://doi.org/10.14302/issn.2578-8590.ipj-18-2524>

Keywords: neurohormone, affective disorders, psychosomatic diseases and cancer

Received: Dec 06, 2018

Accepted: Dec 11, 2018

Published: Dec 18, 2018

Editor: Nasim Habibzadeh, Teesside university, UK.

Background

The pineal gland constitutes a major neuroendocrine organ in the brain. By means of its neurohormone melatonin it transduces exogenous signals such as circadian and seasonal variations of light and temperature into proper hormonal changes which adjust and adapt internal endocrine functions. Alteration of circadian rhythms has been associated with affective disorders, psychosomatic diseases and cancer [1, 2].

It has been observed that light deprivation, which stimulates (the enzymes responsible for) melatonin production in the pineal [3, 4] enhances the animal's ethanol preference [5, 6]. Similarly, administration of the pineal hormone to rats maintained under normal conditions of constant photoperiod also induced ethanol drinking [7 – 9].

The pineal gland synthesizes serotonin and converts it to melatonin [10]. It is then possible to postulate an influence-interaction between the serotonergic system upon pineal activities. Specifically, it has been reported that cocaine given to adult rats stimulated pineal melatonin synthesis through increased 5HT-N-acetyltransferase activity in rat pineal gland [11, 12]. Evidence for selective inhibition of median raphe neuron metabolism by treatment with melatonin and serotonin has been also described with stability of the raphe parameter in presence of noradrenaline (NA), histamine, dopamine (DA) [13, 14].

Several observations have demonstrated that the opiate system can modulate melatonin secretion from the pineal gland and that the effect of opiates may require a pineal participation. In particular, Lissoni et al. [15] reported that treatment with melatonin diminished plasma levels of B-endorphin in men. Acute administration of morphine resulted in a dose dependent increase in plasma melatonin concentration and this effect was blocked by pretreatment with naloxone [16]. Thus, these results seem to support the hypothesis that the opioidergic system might contribute to the activation of melatonin secretion as also proposed more recently [17, 18].

Direct involvement of melatonin within the activity of the dopaminergic system in the nucleus accumbens (nAcc) that is specifically involved in reward [19, 20] has been suggested by Gaffori et

al. [21]. Their data are indeed showing a decreased locomotor activity after injection of melatonin into the nAcc of rats. This action of melatonin was completely antagonized by endorphins, and this introduces the hypothesis of a complex interaction between the DA system, the opiate system and the pineal hormone [16, 22].

In this context may also be involved data showing that treatment with gamma endorphin as well as haloperidol (DA antagonist) were followed by increased levels of melatonin [23]. Ex vivo data indicated a significant increase in brain DA and NA after treatment with melatonin [24, 25] while inhibition of the release of DA in vitro in presence of melatonin has been also reported [26, 27] and this inhibition clearly exhibited a 24-hours rhythm [28, 29].

The Pineal - Melatonin Hypothesis

All these preliminary informations seem to indicate a close interrelationship between melatonin and the "biochemical markers" involved in the addiction - reward - craving states. This seems true for alcohol, as suggested above and by evidence that alcohol withdrawal syndrome produced a reduction of nocturnal pineal melatonin content with a concomitant elevation in pineal serotonin. In addition, a group of "abstinent" (polydrug addicted) showed remarkably higher melatonin levels than acute relapsive cases [30, 31]. It could be possible, in view of these preliminary informations, that melatonin is directly involved in the development of a state of drug addiction.


Our preliminary hypothesis could be that in conditions of drug intake, synthesis and release of melatonin in the pineal is altered (probably abnormally increased), either primarily or following alteration of the other neural systems defined above (DA, 5HT, and/or opiate). In normal conditions melatonin might be acting as a cerebral "pacemaker", sensitive to endogenous as well as exogenous stimuli in the attempt to maintain an equilibrate circadian interaction between the cerebral activities of endogenous aminergic and opiates systems (probably via a circadian mechanism). However, abnormal states (i.e. drug intake, abuse, addiction) could result in chronically altered pineal functions, resulting in rhythmically higher synthesis and release of melatonin (increased turnover). This could be either an

original effect of drug intake or subsequent to modified activity of aminergic (dopaminergic, serotonergic) and/or opioidergic systems following drug abuse [32, 33]. This change in the pineal functions could result in a rhythmically abnormal change in the functions of cerebral opiates (which should be reduced) and/or monoamine neurotransmitters, i.e. altered functions of DA in the nAcc (with a rhythmically higher increase and decrease of DA activity) and/or opposite modification of

serotonergic functions, which could end in the development of a "circadian" state of reward - urge for drug (craving) resulting in addiction (see the resulting scheme 1).

Proposed Strategy to Study the Pineal - Melatonin Hypothesis

This hypothesis could be studied in models of addicted animals i.e. spontaneous alcohol drinking rats



Scheme 1. circadian state of reward - urge for drug (craving) resulting in addiction

versus spontaneous non preferring water drinking rats selected as described earlier [34] then submitted to treatment with melatonin receptor antagonists and/or agonists.

Experimental Procedures

The above mentioned compounds, as well as melatonin, will be injected (intracerebral and/or systemically) in control and addicted animals. If the "melatonin hypothesis" is correct, one would expect modification of ethanol intake in spontaneous alcohol drinking rats. Similarly, modifications should be detectable in the behaviour of addicted animals submitted to the conditioned place preference test as well as to self administration procedures (either i.v. [35, 36] or intracranial self administration of drugs [37, 38]. Modifications of neuro-biochemical parameters such as amine neurotransmitters and/or endogenous enkephalin and endorphin functions and receptor activities should be correlated to these behavioural alterations and monitored by mean of in vitro methods i.e. autoradiography [39] and in vivo techniques such as intracranial microdialysis [40] and in vivo electrochemistry [41]. In particular, previous in vitro and in vivo experiments using the electrochemical method of voltammetry [42] indicated that Melatonin is an electroactive indoleamine hormone and that it can be measurable in vivo in the pineal gland as well as in the suprachiasmatic nucleus of rat brain [43]. This, together with the possibility to concomitant voltammetric monitoring of DA and 5-HT release and metabolism [42, 44] will be a very useful tool to perform such investigation and already preliminary data have proposed a significant influence of melatonin or its antagonism on alcohol consumption in ethanol drinking rats [8]. Further data supporting the proposed hypothesis will be of help in designing new therapeutic approaches to tackle drug dependence.

References

1. Moss, H., Tamarkin, L., Martin, P., & Linnoila, M. (1986). Pineal function during ethanol intoxication, dependence and withdrawal. *Life Sci.*, 39, 2209-2214.
2. Peres, R., do Amaral, F.G., Madrigrano, T.C., Scialfa, J.H., Bordin, S. et al. (2011). Ethanol consumption and pineal melatonin daily profile in rats. *Addiction Biology*, 16, 580-590. <https://doi.org/10.1111/j.1369-1600.2011.00342.x>

3. Axelrod, J., Wurtman, R. & Snyder, S. (1965). Control of hydroxyindole-O-methyltransferase activity in the rat pineal gland by environmental lighting. *J. Biol. Chem.*, 240, 949-954.
4. Munro, H.N. & Allison, J.B. (2014) *Mammalian Protein Metabolism, Volume 1* -Academic Press, 582 pp. books.google.com
5. Reiter, R., Blum, K., Wallace, J. & Merritt, J. (1973). Effect of the pineal gland on alcohol consumption. *Q. J. Stud. Alcohol.* 34, 937-939.
6. Vengeliene, V., Noori, H.R. & Spanagel, R. (2015). Activation of melatonin receptors reduces relapse-like alcohol consumption *Neuropsychopharmacology*, 40, 2897–2906.
7. Geller, I., Purdy, R. & Merritt, J. (1973). Alterations in ethanol preference in the rat : role of brain biogenic amines. *Ann. N.Y. Acad. Sci.* 215, 54-59.
8. Crespi, F. (2012). Influence of melatonin or its antagonism on alcohol consumption in ethanol drinking rats: a behavioral and in vivo voltammetric study *Brain Res*, 1452, 39-46. <https://doi.org/10.1016/j.brainres.2011.10.050>
9. Deehan, Jr. G.A., Hauser, S.R., Wilden, J.A., Truitt, W.A. & Rodd, Z.A. (2013). Elucidating the biological basis for the reinforcing actions of alcohol in the mesolimbic dopamine system: the role of active metabolites of alcohol. *Front. Behav. Neurosci.*, 7, 104. doi.org/10.3389/fnbeh.2013.00104
10. Brzezinski, A. (1997). Melatonin in humans. *N. Engl. J. Med.*, 336, 186-195. DOI: 10.1056/NEJM199701163360306
11. Oxenkrug, G., Dragovic, L., Marks, B. & Yuwiler, A. (1990). Effect of cocaine on rat pineal melatonin synthesis in vivo and in vitro. *Psychiatry Res.*, 34, 185-192.
12. de Mesquita, L.S.M., Garcia, R.C.T. & Amaral, F.G. (2017). The muscarinic effect of anhydroecgonine methyl ester, a crack cocaine pyrolysis product, impairs melatonin synthesis in the rat pineal gland. *Toxicol. Res.*, 6, 420-431. DOI: 10.1039/C7TX00009J
13. Lloyd, O.L. (1974). Inhibition of medial raphe neuron metabolism by cerebro spinal fluid containing

- 5HT and melatonin. *Biochem. Pharmacol.*, 23, 1913-1915.
14. Leite-Panissi, C.R.A., Ferrarese, A.A. & Terzian, A.L.B. (2006). Serotonergic activation of the basolateral amygdala and modulation of tonic immobility in guinea pig *Brain Research Bulletin*, 69, 356-364.
 15. Lissoni, P., Esposti, D., Mauri, R. & Fraschini, F. (1986). A clinical study on the relationship between the pineal gland and the opioid system. *J. Neural Transm.*, 65, 63-73.
 16. Esposti, D., Esposti, G., Lissoni, P. & Fraschini, F. (1988). The pineal gland - opioid system relation: melatonin-naloxone interactions in regulating GH and LH releases in man. *J. Endocrinol. Invest.*, 11, 103-106.
 17. Chuchuen, U., Ebadi, M. & Govitrapong, P. (2004). The stimulatory effect of mu- and delta-opioid receptors on bovine pinealocyte melatonin synthesis *Journal of Pineal Research*, 37, 223-229.
 18. Pačesová, D., Novotný, J. & Bendová, Z. (2016). The Effect of Chronic Morphine or Methadone Exposure and Withdrawal on Clock Gene Expression in the Rat Suprachiasmatic Nucleus and AA-N AT Activity in the Pineal Gland. *Physiological Research*, 65, 517-525. ISSN 1802-9973(online).
 19. O'connell, L.A. & Hofmann, H.A. (2011). The Vertebrate mesolimbic reward system and social behavior network: A comparative synthesis. *Journal of Comparative Neurology*, 519, 3599-3639. <https://doi.org/10.1002/cne.22735>.
 20. Ikemoto, S. & Panksepp, J. (1999). The role of nucleus accumbens dopamine in motivated behavior: a unifying interpretation with special reference to reward-seeking. *Brain Research Reviews*, 31, 6-41. [https://doi.org/10.1016/S0165-0173\(99\)00023-5](https://doi.org/10.1016/S0165-0173(99)00023-5).
 21. Gaffori, O., & Van Ree, JM. (1985). β -Endorphin-(10–16) antagonizes behavioral responses elicited by melatonin following injection into the nucleus accumbens of rats. *Life sciences*, 37, 357-364.
 22. Ibáñez-Costa, A., Córdoba-Chacón, J., Gahete, M.D. Kineman, R.D., Castaño, J.P. & Luque, R.M. (2015). Melatonin Regulates Somatotrope and Lactotrope Function Through Common and Distinct Signaling Pathways in Cultured Primary Pituitary Cells From Female Primates. *Endocrinology*, 156, 1100–1110, <https://doi.org/10.1210/en.2014-1819>.
 23. Gaffori, O., Geffard, M., Van Ree, J. (1983). des-Tyr1-gamma-endorphin and haloperidol increase pineal gland melatonin levels in rats. *Peptides*, 4, 393-395.
 24. Wendel, O., Waterbury, L. & Pearce, L. (1974). Increase in monoamine concentration in rat brain following melatonin administration. *Experientia*, 30, 1167-1168.
 25. Cassim, L., Maharaj, D.S., Maharaj, H. & Daya, S. (2006). Melatonin Counters the 5-fluorouracil-induced Decrease in Brain Serotonin and Dopamine Levels. *Annals of Neurosciences, Chandigarh* 13, Fasc 2, 4pp DOI:10.5214/ans.0972.7531.2006.130201
 26. Zisapel, N. & Laudon, M. (1983). Inhibition by melatonin of dopamine release from rat hypothalamus: regulation of calcium entry. *Brain Res.*, 272, 378-381.
 27. Claustrat, B., Valatx, J.L., Harthe, C. & Brun, J. (2008). Effect of Constant Light on Prolactin and Corticosterone Rhythms Evaluated Using a Noninvasive Urine Sampling Protocol in the Rat. *Horm. Metab. Res.*, 40, 398-403, DOI:10.1055/s-2008-1065330
 28. Zisapel, N., Egozi, Y. & London, M. (1985). Circadian variations in the inhibition of DA release from rat hypothalamus by melatonin. *Neuroendocrinology*, 40, 102-108.
 29. Pandi-Perumal, S.R., Trakht, I., Srinivasan, V., Spence, D.V. & Maestroni, J.M. (2008). Physiological effects of melatonin: Role of melatonin receptors and signal transduction pathways. *Progress in Neurobiology*, 85, 335-353.
 30. Veit, I., Dietzel, M., Lesch, O., Hermann, P., Birsak, L. & Reschenhofer, E. (1986). Circadian neuroendocrinologic profile in patients with multiple drug abuse. *Wien Med. Wochenschr.*, 136, 500-504.
 31. Bondi, C.D., Kamal, K.M., Johnson, D.A., Witt-Enderby, P.A. & Giannetti, V.J. (2018). The Effect of Melatonin Upon Postacute Withdrawal

- Among Males in a Residential Treatment Program (M-PAWS): A Randomized, Double-blind, Placebo-controlled Trial. *Journal of Addiction Medicine*, 2, 201–206. doi: 10.1097/ADM.0000000000000386.
32. Eiden, L.E. & Weihe, E. (2011). VMAT2: a dynamic regulator of brain monoaminergic neuronal function interacting with drugs of abuse. *Annals NYAS*, 1216, 86-98. <https://doi.org/10.1111/j.1749-6632.2010.05906.x>.
 33. Gianoulakis, C. (2004). Endogenous Opioids and Addiction to Alcohol and other Drugs of Abuse. *Current Topics in Medicinal Chemistry*, 4, 39-50. DOI: <https://doi.org/10.2174/1568026043451573>.
 34. Crespi, F. (1998). The role of cholecystokinin (CCK), CCK-A or CCK-B receptor antagonists in the spontaneous preference for drugs of abuse (alcohol or cocaine) in naive rats. *Methods Find. Exp. Clin. Pharmacol.*, 20, (8) 679-697.
 35. Weeks, J. (1962). Experimental morphine addiction: Method for automatic intravenous injections in unrestrained rats. *Science*, 138, 143-144.
 36. Knackstedt ,L.A. & Kalivas, P.W. (2007). Extended Access to Cocaine Self-Administration Enhances Drug-Primed Reinstatement but Not Behavioral Sensitization. *Journal of Pharmacology and Experimental Therapeutics*, 322, 1103-1109. DOI: <https://doi.org/10.1124/jpet.107.122861>.
 37. Goldberg, S.R., Spealman, R.D. & Kelleher, R.T. (1979). Enhancement of drug-seeking behavior by environmental stimuli associated with morphine or cocaine injections. *Neuropharmacology*, 18, 1015-1017.
 38. Carlezon Jr., W.A. & Chartoff, E.H. (2007). Intracranial self-stimulation (ICSS) in rodents to study the neurobiology of motivation. *Nature Protocols*, 2, 2987–2995.
 39. Tempel, A. & Zukin, R.S. (1987). Neuroanatomical patterns of the mu, delta, and kappa opioid receptors of rat brain as determined by quantitative in vitro autoradiography. *PNAS*, 84, 4308-4312. <https://doi.org/10.1073/pnas.84.12.4308>.
 40. Richter, R.M.& Weiss, F. (1999). In vivo CRF release in rat amygdala is increased during cocaine withdrawal in self-administering rats. *Synapse*, 32, 254-261. [https://doi.org/10.1002/\(SICI\)1098-2396\(19990615\)32:4<254::AID-SYN2>3.0.CO;2-H](https://doi.org/10.1002/(SICI)1098-2396(19990615)32:4<254::AID-SYN2>3.0.CO;2-H).
 41. Martin, K.F., Marsden, C.A. & Crespi, F. In vivo electrochemistry with carbon fibre electrodes: principles and application to neuropharmacology. *Trends Anal. Chem.*, 7, 334-339.
 42. Crespi, F. (1986). Differential pulse voltammetry, a new elegant methodology for the in vivo exploration of biogenic amines in specific brain areas. *Medicina, Rivista della Enciclopedia Medica Italiana (Riv E.M.I.)*, 6, 154-158.
 43. Crespi, F., Benedetti, R. & Marien, M. (1992). Melatonin an electroactive indoleamine hormone measurable in vivo by voltammetry. *Neuroscience Letters*, 42, S8.
 44. Crespi, F. (2013). Invasive or Non-Invasive Techniques and Sensors for Real Time In Vivo Sensing in the Brain" chapter in the book, *Laboratory and Clinical Research: "Microelectrodes: Techniques, Structures for Biosensing and Potential Applications"* Nova Science Publishers, Inc.